Author Index

Aşçı, S. *see* Haznedaroğlu, F.
Adams, G.G. *see* Ehrmann, E.H.
Aeinehchi, M., Eslami, B., Ghanbariha, M. & Saffar, A.S. Mineral trioxide aggregate (MTA) and calcium hydroxide as pulp-capping agents in human teeth: a preliminary report, 225
Ahn, S. *see* Martín, B.
Ajam, F. *see* Attin, T.
Al-Fouzan, K.S. Incidence of rotary ProFile instrument fracture and the potential for bypassing *in vivo*, 864
Al-Kilani, M.G., Whitworth, J.M. & Dummer, P.M.H. Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
Al-Yahya, A.S. *see* Saad, A.Y.
Ambu, R. *see* Cotti, E.
Andrade Feitosa Jr, J.P. *see* Gurgel-Filho, E.D.
Andrade, A.F.B. *see* Rôças, I.N.
Angyal, J. *see* Dobó-Nagy, C.
Anić, I. *see* Miletić, I.
Anić, I. *see* Vidučić, D.
Appel, T. *see* Braun, A.
Araújo Silva, J.B. *see* Gurgel-Filho, E.D.
Ardila, C.N. *see* Wu, M.-K.
Ardila, C.N., Wu, M-K. & Wesselink, P.R. Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591
von Arx, T., Montagne, D., Zwinggi, C. & Lussi, A. Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
Asgari, S., Janahmadi, M. & Khalilkhani, H. Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
Attin, T., Paqué, F., Ajam, F. & Lennon, Á.M. Review of the current status of tooth whitening with the walking bleach technique, 313

Bahcall, J. & Barss, J. Orascopic visualization technique for conventional and surgical endodontics, 441
Bahillo, J.G. *see* Martín, B.
Bal, S. *see* Boussetta, F.
Barbakow, F. *see* Hänni, S.
Barbakow, F. *see* Hübscher, W.
Barbakow, F. *see* Peters, O.A.
Barss, J. *see* Bahcall, J.
Batur, B. *see* Haznedaroğlu, F.
Beke, D. *see* Dobó-Nagy, C.
Bergenholtz, G. *see* Chávez de Paz, L.E.
Bergenholtz, G. *see* Ricucci, D.
Bergmans, L., Van Cleynenbreugel, J., Beullens, M., Wevers, M., Van Meerbeek, B. & Lambrechts, P. Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
Bernáth, M. & Szabó, J. Tissue reaction initiated by different sealers, 256
Beullens, M. *see* Bergmans, L.
Bier, C.A. *see* Kopper, P.M.P.
Bilgic, B. *see* Ozbas, H.
Boltacz-Rzepkowska, E. & Pawlicka, H. Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
Božić, Z. *see* Vidučić, D.
Boivin, G. *see* Boussetta, F.
Bonetti Filho, I. *see* Tanomaru, J.M.G.
Bopp, S. *see* Kopper, P.M.P.
Boussetta, F., Bal, S., Romeas, A., Boivin, G., Magloire, H. & Farge, P. *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical gutta-percha condensation techniques, 367
Brückner, M. *see* Weiger, R.
Braem, M. *see* Hommez, G.M.G.
Braun, A., Appel, T. & Frentzen, M. Endodontic and surgical treatment of a geminated maxillary incisor, 380
Breschi, L. *see* Venturi, M.
Briseño Marroquin, B. *see* Pistorius, A.
Brugnera-Junior, A. *see* Picoli, F.
Burhanoğlu, D. *see* Önçağ, Ö.

Campisi, G. *see* Cotti, E.
Canzobre, M.C. *see* Kaplan, A.E.
Capelli, G. *see* Venturi, M.
Caputi, S. *see* Esposito, P.
Castellanos, A.V. *see* Espina, A.I.
Cathro, P.R. & Love, R.M. Comparison of MicroSeal and System B/Obtura II obturation techniques, 876
Çalşkan, G. *see* Çalşkan, M.K.
Çalşkan, M.K., Öztop, F. & Çalşkan, G. Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64
Chávez de Paz, L.E., Dahlén, G., Molander, Å., Möller, A. & Bergenholtz, G. Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
Chan, T.K. *see* Cheung, G.S.P.
Chandler, N.P. *see* Hauman, C.H.J.
Chandler, N.P. *see* Naoum, H.J.

© 2003 Blackwell Publishing Ltd

Author Index

Aşçı, S. *see* Haznedaroğlu, F.
Adams, G.G. *see* Ehrmann, E.H.
Aeinehchi, M., Eslami, B., Ghanbariha, M. & Saffar, A.S. Mineral trioxide aggregate (MTA) and calcium hydroxide as pulp-capping agents in human teeth: a preliminary report, 225
Ahn, S. *see* Martín, B.
Ajam, F. *see* Attin, T.
Al-Fouzan, K.S. Incidence of rotary ProFile instrument fracture and the potential for bypassing *in vivo*, 864
Al-Kilani, M.G., Whitworth, J.M. & Dummer, P.M.H. Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
Al-Yahya, A.S. *see* Saad, A.Y.
Ambu, R. *see* Cotti, E.
Andrade Feitosa Jr, J.P. *see* Gurgel-Filho, E.D.
Andrade, A.F.B. *see* Rôças, I.N.
Angyal, J. *see* Dobó-Nagy, C.
Anić, I. *see* Miletić, I.
Anić, I. *see* Vidučić, D.
Appel, T. *see* Braun, A.
Araújo Silva, J.B. *see* Gurgel-Filho, E.D.
Ardila, C.N. *see* Wu, M.-K.
Ardila, C.N., Wu, M-K. & Wesselink, P.R. Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591
von Arx, T., Montagne, D., Zwinggi, C. & Lussi, A. Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
Asgari, S., Janahmadi, M. & Khalilkhani, H. Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
Attin, T., Paqué, F., Ajam, F. & Lennon, Á.M. Review of the current status of tooth whitening with the walking bleach technique, 313

Bahcall, J. & Barss, J. Orascopic visualization technique for conventional and surgical endodontics, 441
Bahillo, J.G. *see* Martín, B.
Bal, S. *see* Boussetta, F.
Barbakow, F. *see* Hänni, S.
Barbakow, F. *see* Hübscher, W.
Barbakow, F. *see* Peters, O.A.
Barss, J. *see* Bahcall, J.
Batur, B. *see* Haznedaroğlu, F.
Beke, D. *see* Dobó-Nagy, C.
Bergenholtz, G. *see* Chávez de Paz, L.E.
Bergenholtz, G. *see* Ricucci, D.
Bergmans, L., Van Cleynenbreugel, J., Beullens, M., Wevers, M., Van Meerbeek, B. & Lambrechts, P. Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
Bernáth, M. & Szabó, J. Tissue reaction initiated by different sealers, 256
Beullens, M. *see* Bergmans, L.
Bier, C.A. *see* Kopper, P.M.P.
Bilgic, B. *see* Ozbas, H.
Boltacz-Rzepkowska, E. & Pawlicka, H. Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
Božić, Z. *see* Vidučić, D.
Boivin, G. *see* Boussetta, F.
Bonetti Filho, I. *see* Tanomaru, J.M.G.
Bopp, S. *see* Kopper, P.M.P.
Boussetta, F., Bal, S., Romeas, A., Boivin, G., Magloire, H. & Farge, P. *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical gutta-percha condensation techniques, 367
Brückner, M. *see* Weiger, R.
Braem, M. *see* Hommez, G.M.G.
Braun, A., Appel, T. & Frentzen, M. Endodontic and surgical treatment of a geminated maxillary incisor, 380
Breschi, L. *see* Venturi, M.
Briseño Marroquin, B. *see* Pistorius, A.
Brugnera-Junior, A. *see* Picoli, F.
Burhanoğlu, D. *see* Önçağ, Ö.

Campisi, G. *see* Cotti, E.
Canzobre, M.C. *see* Kaplan, A.E.
Capelli, G. *see* Venturi, M.
Caputi, S. *see* Esposito, P.
Castellanos, A.V. *see* Espina, A.I.
Cathro, P.R. & Love, R.M. Comparison of MicroSeal and System B/Obtura II obturation techniques, 876
Çalşkan, G. *see* Çalşkan, M.K.
Çalşkan, M.K., Öztop, F. & Çalşkan, G. Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64
Chávez de Paz, L.E., Dahlén, G., Molander, Å., Möller, A. & Bergenholtz, G. Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
Chan, T.K. *see* Cheung, G.S.P.
Chandler, N.P. *see* Hauman, C.H.J.
Chandler, N.P. *see* Naoum, H.J.

© 2003 Blackwell Publishing Ltd

Chandler, N.P., Pitt Ford, T.R. & Monteith, B.D. Coronal pulp size in molars: a study of bitewing radiographs, 757
Chang, Y.-C. *see* Yang, L.-C.
Chen, C.-S. *see* Chueh, L.-H.
Chen, S.-C. *see* Chueh, L.-H.
Cheung, G.S.P. & Chan, T.K. Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
Choi, M.Y. *see* Lim, T.S.
Chong, B.S., Pitt Ford, T.R. & Hudson, M.B. A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
Chueh, L.-H., Chen, S.-C., Lee, C.-M., Hsu, Y.-Y., Pai, S.-F., Kuo, M.-L., Chen, C.-S., Duh, B.-R., Yang, S.-F., Tung, Y.-L. & Hsiao, C.K. Technical quality of root canal treatment in Taiwan, 416
Costa, C.A.S., Oliveira, M.F., Giro, E.M.A. & Hebling, J. Biocompatibility of resin-based materials used as pulp-capping agents, 831
Cotti, E., Campisi, G., Ambu, R. & Dettori, C. Ultrasound real-time imaging in the differential diagnosis of periapical lesions, 556
Cox, C.F. *see* Murray, P.E.

Dahlén, G. *see* Chávez de Paz, L.E.
Daróczi, L. *see* Dobó-Nagy, C.
De Moor, R.J.G. *see* Hommez, G.M.G.
Della Bona, A. *see* Kopper, P.M.P.
Delschen, S. *see* Sonntag, D.
Dettori, C. *see* Cotti, E.
Dobó-Nagy, C., Fejérdy, P., Angyal, J., Harasztosi, L., Daróczi, L., Beke, D. & Wesselink, P.R. Measurement of periapical pressure created by occlusal loading, 700
Dorn, S. *see* Garala, M.
Dugas, N.N., Lawrence, H.P., Teplitsky, P.E., Pharoah, M.J. & Friedman, S. Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations, 181
Duh, B.-R. *see* Chueh, L.-H.
Dummer, P.M.H. *see* Al-Kilani, M.G.

Ebner, J.P. *see* Velvart, P.
Ebner-Zimmermann, U. *see* Velvart, P.
Economides, N., Pantelidou, O., Kokkas, A. & Tziafas, D. Short-term periradicular tissue response to mineral trioxide aggregate (MTA) as root-end filling material, 44
Ehrmann, E.H., Messer, H.H. & Adams, G.G. The relationship of intracanal medicaments to postoperative pain in endodontics, 868
ElAyouti, A. *see* Weiger, R.
Er, K. *see* Hürmüzlü, F.
Eronat, C. *see* Önçağ, Ö.
Ersev, H. *see* Haznedaroğlu, F.
Eslami, B. *see* Aeinehchi, M.

Espina, A.I., Castellanos, A.V. & Fereira, J.L. Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
Esposito, P., Varvara, G., Caputi, S. & Perinetti, G. Catalase activity in human healthy and inflamed dental pulps, 599

Falconi, M. *see* Venturi, M.
Farge, P. *see* Boussetta, F.
Fejérdy, P. *see* Dobó-Nagy, C.
Fenno, J.C. *see* Shur, A.L.
Fereira, J.L. *see* Espina, A.I.
Ferraz, C.C.R. *see* Gomes, B.P.F.A.
Ferraz, C.C.R. *see* Gondim Jr, E.
Ferraz, C.C.R. *see* Pinheiro, E.T.
Ferraz, C.C.R. *see* Teixeira, F.B.
Fidel, R. *see* Sassone, L.M.
Fidel, S. *see* Sassone, L.M.
Figueiredo, J.A.P. *see* Kopper, P.M.P.
Florek, H. *see* Schäfer, E.
Foxton, R.M. *see* Hiraishi, N.
Frentzen, M. *see* Braun, A.
Friedman, S. *see* Dugas, N.N.
Fuss, Z. *see* Kfir, A.

Gagliani, M. *see* Maddalone, M.
Galgano, C., Samson, J., Küffer, R. & Lombardi, T. Focal cemento-osseous dysplasia involving a mandibular lateral incisor, 907
Garaj-Vrhovac, V. *see* Miletić, I.
Garala, M., Kuttler, S., Hardigan, P., Steiner-Carmi, R. & Dorn, S. A comparison of the minimum canal wall thickness remaining following preparation using two nickel–titanium rotary systems, 636
Garcia, J.M.Q., Martins, M.D., Jaeger, R.G. & Marques, M.M. Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404
Ghanbariha, M. *see* Aeinehchi, M.
Ghossayn, F. *see* Yared, G.
Giro, E.M.A. *see* Costa, C.A.S.
Gomes, B.P.F.A. *see* Gondim Jr, E.
Gomes, B.P.F.A. *see* Pinheiro, E.T.
Gomes, B.P.F.A. *see* Teixeira, F.B.
Gomes, B.P.F.A., Sato, E., Ferraz, C.C.R., Teixeira, F.B., Zaia, A.A. & Souza-Filho, F.J. Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
Gomes, B.P.F.A., Souza, S.F.C., Ferraz, C.C.R., Teixeira, F.B., Zaia, A.A., Valdrighi, L. & Souza-Filho, F.J. Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267
Gomez, R.S. *see* Pimenta, F.J.G.S.
Gondim Jr, E., Zaia, A.A., Gomes, B.P.F.A., Ferraz, C.C.R., Teixeira, F.B. & Souza-Filho, F.J. Investigation of the

© 2003 Blackwell Publishing Ltd

marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491
Grawehr, M., Sener, B., Waltimo, T. & Zehnder, M. Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
Gressmann, G. *see* Hülsmann, M.
Guerisoli, D.M.Z. *see* Picoli, F.
Gulabivala, K. *see* McMichen, F.R.S.
Guntermann, A. *see* Sonntag, D.
Gurgel-Filho, E.D., Andrade Feitosa, J.P., Teixeira, F.B., Monteiro de Paula, R.C., Araújo Silva Jr, J.B. & Souza-Filho, F.J. Chemical and X-ray analyses of five brands of dental gutta-percha cone, 302
Gutmann, J.L. *see* Pérez, A.L.
Gutmann, J.L. *see* Satchell, P.G.

Hänni, S., Schönenberger, K., Peters, O.A. & Barbakow, F. Teaching an engine-driven preparation technique to undergraduates: initial observations, 476
Hübscher, W., Barbakow, F. & Peters, O.A. Root canal preparation with FlexMaster: asessment of torque and force in relation to canal anatomy, 883
Hübscher, W., Barbakow, F. & Peters, O.A. Root-canal preparation with FlexMaster: canal shapes analysed by micro-computed tomography, 740
Hülsmann, M. *see* Rödig, T.
Hülsmann, M., Gressmann, G. & Schäfers, F. A comparative study of root canal preparation using FlexMaster and HERO 642 rotary Ni–Ti instruments, 358
Hülsmann, M., Heckendorff, M. & Lennon, Á. Chelating agents in root canal treatment: mode of action and indications for their use, 810
Hülsmann, M., Herbst, U. & Schäfers, F. Comparative study of root-canal preparation using Lightspeed and Quantec SC rotary NiTi instruments, 748
Hürmüzlü, F., Serper, A., Siso, Ş.H. & Er, K. *In vitro* fracture resistance of root-filled teeth using new-generation dentine bonding adhesives, 770
Haenni, S., Schmidlin, P.R., Mueller, B., Sener, B. & Zehnder, M. Chemical and antimicrobial properties of calcium hydroxide mixed with irrigating solutions, 100
Hafez, A.A. *see* Murray, P.E.
Harasztosi, L. *see* Dobó-Nagy, C.
Hardigan, P. *see* Garala, M.
Hauman, C.H.J., Chandler, N.P. & Purton, D.G. Factors influencing the removal of posts, 687
Hauman, C.H.J. & Love, R.M. Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 1. Intracanal drugs and substances, 75
Hauman, C.H.J. & Love, R.M. Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 2. Root-canal-filling materials, 147
Haznedaroğlu, F., Ersev, H., Odabaşı, H., Yetkin, G., Batur, B., Aşçı, S. & İşsever, H. Incidence of patent furcal accessory canals in permanent molars of a Turkish population, 515
Hebling, J. *see* Costa, C.A.S.
Heckendorff, M. *see* Hülsmann, M.
Herbison, P. *see* Naoum, H.J.
Herbst, U. *see* Hülsmann, M.
Hilmioğlu, S. *see* Önçağ, Ö.
Hiraishi, N., Kitasako, Y., Nikaido, T., Foxton, R.M., Tagami, J. & Nomura, S. Acidity of conventional luting cements and their diffusion through bovine dentine, 622
Hirata, R. *see* Sassone, L.M.
Hoşgör, M. *see* Önçağ, Ö.
Hommez, G.M.G., Braem, M. & De Moor, R.J.G. Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
Hommez, G.M.G., De Moor, R.J.G. & Braem, M. Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
Hsiao, C.K. *see* Chueh, L.-H.
Hsu, Y.-Y. *see* Chueh, L.-H.
Hsu, Y.-Y., Jou, Y.-T., Wong, R., Karabucak, B., Simchon, S. & Kim, S. Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
Huang, F.-M. *see* Yang, L.-C.
Hudson, M.B. *see* Chong, B.S.
Huumonen, S., Lenander-Lumikari, M., Sigurdsson, A. & Ørstavik, D. Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide–eugenol-based sealer, 296

İşsever, H. *see* Haznedaroğlu, F.
Ikeda, H. *see* Puapichartdumrong, P.
Ishikawa, H., Sawada, N., Kobayashi, C. & Suda, H. Evaluation of root-end cavity preparation using ultrasonic retrotips, 586
Issever, H. *see* Ozbas, H.
Järvensivu, A. *see* Waltimo, T.
Jaeger, R.G. *see* Garcia, J.M.Q.
Janahmadi, M. *see* Asgari, S.
Jou, Y.-T. *see* Hsu, Y.-Y.
Jukić, S. *see* Miletić, I.
Jukić, S. *see* Vidučić, D.

Küffer, R. *see* Galgano, C.
Kaplan, A.E., Ormaechea, M.F., Picca, M., Canzobre, M.C. & Ubios, A.M. Rheological properties and biocompatibility of endodontic sealers, 527
Karabucak, B. *see* Hsu, Y.-Y.
Karlović, Z. *see* Vidučić, D.
Katz, A. & Tamse, A. A combined radiographic and computerized scanning method to evaluate remaining dentine thickness in mandibular incisors after various intracanal procedures, 682
Kaufman, A.Y. *see* Tsesis, I.
Kerezoudis, N.P., Siskos, G.J. & Tsatsas, V. Bilateral buccal radicular groove in maxillary incisors: case report, 898
Kfir, A., Rosenberg, E., Zuckerman, O., Tamse, A. & Fuss, Z. Comparison of procedural errors resulting during

© 2003 Blackwell Publishing Ltd

root canal preparations completed by junior dental students in patients using an '8-step method' versus 'serial step-back technique', 49
Khalilkhani, H. *see* Asgari, S.
Kim, S. *see* Hsu, Y.-Y.
Kim, S.K. *see* Sonntag, D.
Kitasako, Y. *see* Hiraishi, N.
Kobayashi, C. *see* Ishikawa, H.
Koh, W.C. *see* Lim, T.S.
Kokkas, A. *see* Economides, N.
Kopper, P.M.P., Figueiredo, J.A.P., Della Bona, A., Vanni, J.R., Bier, C.A. & Bopp, S. Comparative *in vivo* analysis of the sealing ability of three endodontic sealers in post-prepared root canals, 857
Kulkarni, G.K. *see* Yared, G.
Kuno, T. *see* Tsurumachi, T.
Kuo, M.-L. *see* Chueh, L.-H.
Kuttler, S. *see* Garala, M.
Kuusinen, M. *see* Waltimo, T.

Löst, C. *see* Weiger, R.
Lai, C.-C. *see* Yang, L.-C.
Lambrechts, P. *see* Bergmans, L.
Lawrence, H.P. *see* Dugas, N.N.
Lee, C.-M. *see* Chueh, L.-H.
Lenander-Lumikari, M. *see* Huumonen, S.
Lennon, Á. *see* Hülsmann, M.
Lennon, Á.M. *see* Attin, T.
Leonardo, M.R. *see* Tanomaru, J.M.G.
Leonardo, M.R. *see* Yamashita, J.C.
Leonardo, R.T. *see* Soares, J.A.
Liapatas, S., Nakou, M. & Rontogianni, D. Inflammatory infiltrate of chronic periradicular lesions: an immunohistochemical study, 464
Lim, T.S., Wee, T.Y., Choi, M.Y., Koh, W.C. & Sae-Lim, V. Light and scanning electron microscopic evaluation of Glyde™ File Prep in smear layer removal, 336
Liu, C.-M. *see* Yang, L.-C.
Lombardi, T. *see* Galgano, C.
Love, R.M. *see* Cathro, P.R.
Love, R.M. *see* Hauman, C.H.J.
Love, R.M. *see* Naoum, H.J.
Lussi, A. *see* von Arx, T.

Möller, Å. *see* Chávez de Paz, L.E.
Maddalone, M. & Gagliani, M. Periapical endodontic surgery: a 3-year follow-up study, 193
Magán, F. *see* Martín, B.
Magloire, H. *see* Boussetta, F.
Mandel, E. *see* Mesgouez, C.
Marques, M.M. *see* Garcia, J.M.Q.
Martín, B., Zelada, G., Varela, P., Bahillo, J.G., Magán, F., Ahn, S. & Rodríguez, C. Factors influencing the fracture of nickel-titanium rotary instruments, 262
Martins, M.D. *see* Garcia, J.M.Q.
Matossian, L. *see* Mesgouez, C.
McMichen, F.R.S., Pearson, G., Rahbaran, S. & Gulabivala, K. A comparative study of selected physical properties of five root-canal sealers, 629
Mesgouez, C., Rilliard, F., Matossian, L., Nassiri, K. & Mandel, E. Influence of operator experience on canal preparation time when using the rotary Ni-Ti ProFile system in simulated curved canals, 161
Messer, H.H. *see* Ehrmann, E.H.
Messer, H.H. *see* O'Hoy, P.Y.Z.
Miletić, I. *see* Vidučić, D.
Miletić, I., Jukić, S., Anić, I., Željezić, D., Garaj-Vrhovac, V. & Osmak, M. Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
Mjör, I.A. *see* Murray, P.E.
Molander, A. *see* Chávez de Paz, L.E.
Montagne, D. *see* von Arx, T.
Monteiro de Paula, R.C. *see* Gurgel-Filho, E.D.
Monteith, B.D. *see* Chandler, N.P.
Mueller, B. *see* Haenni, S.
Murray, P.E., Hafez, A.A., Smith, A.J., Windsor, L.J. & Cox, C.F. Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106
Murray, P.E., Smith, A.J., Windsor, L.J. & Mjör, I.A. Remaining dentine thickness and human pulp responses, 33

Nakou, M. *see* Liapatas, S.
Naoum, H.J., Love, R.M., Chandler, N.P. & Herbison, P. Effect of X-ray beam angulation and intraradicular contrast medium on radiographic interpretation of lower first molar root canal anatomy, 12
Nassiri, K. *see* Mesgouez, C.
Niederman, R. & Theodosopoulou, J.N. A systematic review of *in vivo* retrograde obturation materials, 577
Nikaido, T. *see* Hiraishi, N.
Nomura, S. *see* Hiraishi, N.
Nyberg, P. *see* Waltimo, T.

Ørstavik, D. *see* Huumonen, S.
Önçağ, Ö., Hoşgör, M., Hilmioğlu, S., Zekioğlu, O., Eronat, C. & Burhanoğlu, D. Comparison of antibacterial and toxic effects of various root canal irrigants, 423
Öztop, F. *see* Çalşkan, M.K.
O'Hoy, P.Y.Z., Messer, H.H. & Palamara, J.E.A. The effect of cleaning procedures on fracture properties and corrosion of NiTi files, 724
Odabaşı, H. *see* Haznedaroğlu, F.
Oliveira, M.F. *see* Costa, C.A.S.
Opperman, L.A. *see* Pérez, A.L.
Ormaechea, M.F. *see* Kaplan, A.E.
Osmak, M. *see* Miletić, I.
Ozbas, H., Yaltirik, M., Bilgic, B. & Issever, H. Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281

© 2003 Blackwell Publishing Ltd

Pérez, A.L., Spears, R., Gutmann, J.L. & Opperman, L.A. Osteoblasts and MG-63 osteosarcoma cells behave differently when in contact with ProRoot™ MTA and White MTA, 564
Pai, S.-F. *see* Chueh, L.-H.
Palamara, J.E.A. *see* O'Hoy, P.Y.Z.
Pantelidou, O. *see* Economides, N.
Paqué, F. *see* Attin, T.
Pawlicka, H. *see* Boltacz-Rzepkowska, E.
Pearson, G. *see* McMichen, F.R.S.
Pecora, J.D. *see* Picoli, F.
Perinetti, G. *see* Esposito, P.
Peters, C.I. *see* Peters, O.A.
Peters, O.A. *see* Hänni, S.
Peters, O.A. *see* Hübscher, W.
Peters, O.A., Peters, C.I., Schönenberger, K. & Barbakow, F. ProTaper rotary root canal preparation: assessment of torque and force in relation to canal anatomy, 93
Peters, O.A., Peters, C.I., Schönenberger, K. & Barbakow, F. ProTaper rotary root canal preparation: effects of canal anatomy on final shape analysed by micro CT, 86
Pharoah, M.J. *see* Dugas, N.N.
Picca, M. *see* Kaplan, A.E.
Picoli, F., Brugnera-Junior, A., Saquy, P.C., Guerisoli, D.M.Z. & Pecora, J.D. Effect of Er:YAG laser and EDTAC on the adhesiveness to dentine of different sealers containing calcium hydroxide, 472
Pimenta, F.J.G.S., Sá, A.R. & Gomez, R.S. Lymphangiogenesis in human dental pulp, 853
Pinheiro, E.T., Gomes, B.P.F.A., Ferraz, C.C.R., Sousa, E.L.R., Teixeira, F.B. & Souza-Filho, F.J. Microorganisms from canals of root-filled teeth with periapical lesions, 1
Pistorius, A., Willershausen, B. & Briseño Marroquin, B. Effect of apical root-end filling materials on gingival fibroblasts, 610
Pitt Ford, T.R. *see* Chandler, N.P.
Pitt Ford, T.R. *see* Chong, B.S.
Prati, C. *see* Venturi, M.
Puapichartdumrong, P., Ikeda, H. & Suda, H. Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
Purton, D.G. *see* Hauman, C.H.J.

Rôças, I.N. *see* Siqueira Jr, J.F.
Rôças, I.N., Siqueira Jr, J.F., Andrade, A.F.B. & Uzeda, M. Oral treponemes in primary root canal infections as detected by nested PCR, 20
Rödig, T. & Hülsmann, M. Diagnosis and root canal treatment of a mandibular second premolar with three root canals, 912
Rahbaran, S. *see* McMichen, F.R.S.
Ree, M.H., Timmerman, M.F. & Wesselink, P.R. An evaluation of the usefulness of two endodontic case assessment forms by general dentists, 545
Ree, M.H., Timmerman, M.F. & Wesselink, P.R. Factors influencing referral for specialist endodontic treatment amongst a group of Dutch general practitioners, 129
Richardson, M. *see* Waltimo, T.
Ricucci, D. & Bergenholtz, G. Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787
Rilliard, F. *see* Mesgouez, C.
Rodríguez, C. *see* Martín, B.
Romeas, A. *see* Boussetta, F.
Rontogianni, D. *see* Liapatas, S.
Rosenberg, E. *see* Kfir, A.
Rosenberg, E. *see* Tsesis, I.
Rossi, M.A. *see* Yamashita, J.C.

Sá, A.R. *see* Pimenta, F.J.G.S.
Saad, A.Y. & Al-Yahya, A.S. The location of the cementodentinal junction in single-rooted mandibular first premolars from Egyptian and Saudi patients: a histological study, 541
Sae-Lim, V. *see* Lim, T.S.
Saffar, A.S. *see* Aeinehchi, M.
Salo, T. *see* Waltimo, T.
Samson, J. *see* Galgano, C.
Sano, C.L. *see* Teixeira, F.B.
Saquy, P.C. *see* Picoli, F.
Sassone, L.M., Fidel, R., Fidel, S., Vieira, M. & Hirata Jr, R. The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine *in vitro*, 848
Satchell, P.G., Gutmann, J.L. & Witherspoon, D.E. Apoptosis: an introduction for the endodontist, 237
Sato, E. *see* Gomes, B.P.F.A.
Sawada, N. *see* Ishikawa, H.
Schäfer, E. & Florek, H. Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
Schäfer, E. & Schlingemann, R. Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
Schäfer, E. & Zandbiglari, T. Solubility of root-canal sealers in water and artificial saliva, 660
Schäfers, F. *see* Hülsmann, M.
Schönenberger, K. *see* Hänni, S.
Schönenberger, K. *see* Peters, O.A.
Schlingemann, R. *see* Schäfer, E.
Schmidlin, P.R. *see* Haenni, S.
Sedgley, C.M. *see* Shur, A.L.
Sedgley, C.M. & Wagner, R. Orthograde retreatment and apexification after unsuccessful endodontic treatment, retreatment and apicectomy, 780
Sener, B. *see* Grawehr, M.
Sener, B. *see* Haenni, S.
Serper, A. *see* Hürmüzlü, F.

© 2003 Blackwell Publishing Ltd

Shur, A.L., Sedgley, C.M. & Fenno, J.C. The antimicrobial efficacy of 'MGP' gutta-percha *in vitro*, 616
Sigurdsson, A. *see* Huumonen, S.
Silva, L.A.B. *see* Tanomaru, J.M.G.
Silva, L.A.B. *see* Yamashita, J.C.
Simchon, S. *see* Hsu, Y.-Y.
Siqueira Jr, J.F. *see* Rôças, I.N.
Siqueira Jr, J.F. Microbial causes of endodontic flare-ups, 453
Siqueira Jr, J.F. & Rôças, I.N. *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections, 174
Siskos, G.J. *see* Kerezoudis, N.P.
Siso, Ş.H. *see* Hürmüzlü, F.
Sluis, L.W.M. *see* Wu, M.-K.
van der Sluis, L.W.M. *see* Wu, M.-K.
Smith, A.J. *see* Murray, P.E.
Soares, J.A. & Leonardo, R.T. Root canal treatment of three-rooted maxillary first and second premolars – a case report, 705
Sonntag, D., Delschen, S. & Stachniss, V. Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
Sonntag, D., Guntermann, A., Kim, S.K. & Stachniss, V. Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
Sousa, E.L.R. *see* Pinheiro, E.T.
Souza, S.F.C. *see* Gomes, B.P.F.A.
Souza-Filho, F.J. *see* Gomes, B.P.F.A.
Souza-Filho, F.J. *see* Gondim Jr, E.
Souza-Filho, F.J. *see* Gurgel-Filho, E.D.
Souza-Filho, F.J. *see* Pinheiro, E.T.
Souza-Filho, F.J. *see* Teixeira, F.B.
Spears, R. *see* Pérez, A.L.
Stachniss, V. *see* Sonntag, D.
Steinbock, N. *see* Tsesis, I.
Steiner-Carmi, R. *see* Garala, M.
Suda, H. *see* Ishikawa, H.
Suda, H. *see* Puapichartdumrong, P.
Szabó, J. *see* Bernáth, M.

Tagami, J. *see* Hiraishi, N.
Tamse, A. *see* Katz, A.
Tamse, A. *see* Kfir, A.
Tanomaru Filho, M. *see* Tanomaru, J.M.G.
Tanomaru Filho, M. *see* Yamashita, J.C.
Tanomaru, J.M.G., Leonardo, M.R., Tanomaru Filho, M., Bonetti Filho, I. & Silva, L.A.B. Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
Teixeira, F.B. *see* Gomes, B.P.F.A.
Teixeira, F.B. *see* Gondim Jr, E.
Teixeira, F.B. *see* Gurgel-Filho, E.D.
Teixeira, F.B. *see* Pinheiro, E.T.
Teixeira, F.B., Sano, C.L., Gomes, B.P.F.A., Zaia, A.A., Ferraz, C.C.R. & Souza-Filho, F.J. A preliminary *in vitro* study of the incidence and position of the root canal isthmus in maxillary and mandibular first molars, 276
Teplitsky, P.E. *see* Dugas, N.N.
Theodosopoulou, J.N. *see* Niederman, R.
Timmerman, M.F. *see* Ree, M.H.
Tjäderhane, L. *see* Waltimo, T.
Tsai, C.-H. *see* Yang, L.-C.
Tsatsas, V. *see* Kerezoudis, N.P.
Tsesis, I., Steinbock, N., Rosenberg, E. & Kaufman, A.Y. Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372
Tsurumachi, T. & Kuno, T. Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
Tung, Y.-L. *see* Chueh, L.-H.
Tziafas, D. *see* Economides, N.

Ubios, A.M. *see* Kaplan, A.E.
Uzeda, M. *see* Rôças, I.N.

Väänänen, A. *see* Waltimo, T.
Valdrighi, L. *see* Gomes, B.P.F.A.
Van Cleynenbreugel, J. *see* Bergmans, L.
Van Meerbeek, B. *see* Bergmans, L.
Vanni, J.R. *see* Kopper, P.M.P.
Varela, P. *see* Martín, B.
Varvara, G. *see* Esposito, P.
Velvart, P., Ebner-Zimmermann, U. & Ebner, J.P. Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
Venturi, M., Prati, C., Capelli, G., Falconi, M. & Breschi, L. A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
Vidučić, D., Jukić, S., Karlović, Z., Božić, Z., Miletić, I. & Anić, I. Removal of gutta-percha from root canals using an Nd:YAG laser, 670
Vieira, M. *see* Sassone, L.M.

Wagner, R. *see* Sedgley, C.M.
Waltimo, T. *see* Grawehr, M.
Waltimo, T., Kuusinen, M., Järvensivu, A., Nyberg, P., Väänänen, A., Richardson, M., Salo, T. & Tjäderhane, L. Examination on *Candida* spp. in refractory periapical granulomas, 643
Wee, T.Y. *see* Lim, T.S.
Weiger, R., Brückner, M., ElAyouti, A. & Löst, C. Preparation of curved root canals with rotary FlexMaster instruments compared to Lightspeed instruments and NiTi hand files, 483
Wesselink, P.R. *see* Ardila, C.N.
Wesselink, P.R. *see* Dobó-Nagy, C.
Wesselink, P.R. *see* Ree, M.H.
Wesselink, P.R. *see* Wu, M.-K.
Wevers, M. *see* Bergmans, L.
Whitworth, J. Visual Endodontics & Traumatology, 960
Whitworth, J.M. *see* Al-Kilani, M.G.

© 2003 Blackwell Publishing Ltd

Willershausen, B. *see* Pistorius, A.
Windsor, L.J. *see* Murray, P.E.
Witherspoon, D.E. *see* Satchell, P.G.
Wong, R. *see* Hsu, Y.-Y.
Wu, M-K. *see* Ardila, C.N.
Wu, M.-K., van der Sluis, L.W.M., Ardila, C.N. & Wesselink, P.R. Fluid movement along the coronal two-thirds of root fillings placed by three different gutta-percha techniques, 533
Wu, M.-K., Sluis, L.W.M. & Wesselink, P.R. The capability of two hand instrumentation techniques to remove the inner layer of dentine in oval canals, 218

Yaltirik, M. *see* Ozbas, H.
Yamashita, J.C., Tanomaru Filho, M., Leonardo, M.R., Rossi, M.A. & Silva, L.A.B. Scanning electron microscopic study of the cleaning ability of chlorhexidine as a root-canal irrigant, 391
Yang, L.-C., Huang, F.-M., Lin, C.-S., Liu, C.-M., Lai, C.-C. & Chang, Y.-C. Induction of interleukin-8 gene expression by black-pigmented *Bacteroides* in human pulp fibroblasts and osteoblasts, 774
Yang, L.-C., Tsai, C.-H., Huang, F.-M., Liu, C.-M., Lai, C.-C. & Chang, Y.-C. Induction of interleukin-6 gene expression by pro-inflammatory cytokines and black-pigmented *Bacteroides* in human pulp cell cultures, 352
Yang, S.-F. *see* Chueh, L.-H.
Yared, G., Kulkarni, G.K. & Ghossayn, F. An *in vitro* study of the torsional properties of new and used K3 instruments, 764
Yetkin, G. *see* Haznedaroğlu, F.

Željezić, D. *see* Miletić, I.
Zaia, A.A. *see* Gomes, B.P.F.A.
Zaia, A.A. *see* Gondim Jr, E.
Zaia, A.A. *see* Teixeira, F.B.
Zandbiglari, T. *see* Schäfer, E.
Zehnder, M. *see* Grawehr, M.
Zehnder, M. *see* Haenni, S.
Zekioğlu, O. *see* Önçağ, Ö.
Zelada, G. *see* Martin, B.
Zuckerman, O. *see* Kfir, A.
Zwinggi, C. *see* von Arx, T.

© 2003 Blackwell Publishing Ltd

Keyword Index

accessory canals, A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
adhesiveness, Effect of Er:YAG laser and EDTAC on the adhesiveness to dentine of different sealers containing calcium hydroxide, 472
ageing, Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
AH Plus, Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
AH26, Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
amalgam, Effect of apical root-end filling materials on gingival fibroblasts, 610
Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281
anatomical variations, Diagnosis and root canal treatment of a mandibular second premolar with three root canals, 912
Root canal treatment of three-rooted maxillary first and second premolars – a case report, 705
anatomy, A preliminary *in vitro* study of the incidence and position of the root canal isthmus in maxillary and mandibular first molars, 276
angle at fracture, An *in vitro* study of the torsional properties of new and used K3 instruments, 764
antibacterial effects, Comparison of antibacterial and toxic effects of various root canal irrigants, 423
antimicrobial activity, Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
antimicrobial effect, Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
antimicrobial efficacy, Chemical and antimicrobial properties of calcium hydroxide mixed with irrigating solutions, 100
antimicrobial, The antimicrobial efficacy of 'MGP' guttapercha *in vitro*, 616
The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine *in vitro*, 848
apexification, Orthograde retreatment and apexification after unsuccessful endodontic treatment, retreatment and apicectomy, 780
apical leakage, *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical guttapercha condensation techniques, 367
apical periodontitis, Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide–eugenol-based sealer, 296
Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations, 181
apical pressure, Measurement of periapical pressure created by occlusal loading, 700
apical surgery, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
apicectomy, A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
Orthograde retreatment and apexification after unsuccessful endodontic treatment, retreatment and apicectomy, 780
Periapical endodontic surgery: a 3-year follow-up study, 193
apoptosis, Apoptosis: an introduction for the endodontist, 237
automated root canal preparation, A comparative study of root canal preparation using FlexMaster and HERO 642 rotary Ni–Ti instruments, 358
automated root-canal preparation, Comparative study of root-canal preparation using Lightspeed and Quantec SC rotary NiTi instruments, 748

bacteria, Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267
bacterial leakage, Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106
bacterial resistance, Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
Belgium, Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
beta-tricalcium phosphate, Endodontic and surgical treatment of a geminated maxillary incisor, 380
biocompatibility, Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 1. Intracanal drugs and substances, 75
Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 2. Root-canal-filling materials, 147
Biocompatibility of resin-based materials used as pulp-capping agents, 831
Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281

© 2003 Blackwell Publishing Ltd

Rheological properties and biocompatibility of endodontic sealers, 527
biomaterials, Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 1. Intracanal drugs and substances, 75
Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 2. Root-canal-filling materials, 147
Factors influencing the removal of posts, 687
bitewing radiographs, Coronal pulp size in molars: a study of bitewing radiographs, 757
black-pigmented *Bacteroides*, Induction of interleukin-6 gene expression by pro-inflammatory cytokines and black-pigmented *Bacteroides* in human pulp cell cultures, 352
Induction of interleukin-8 gene expression by black-pigmented *Bacteroides* in human pulp fibroblasts and osteoblasts, 774
bleaching, Review of the current status of tooth whitening with the walking bleach technique, 313
blood capillaries, Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
bonding agent, Biocompatibility of resin-based materials used as pulp-capping agents, 831
bone sialoprotein, Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404
bovine dentine, Acidity of conventional luting cements and their diffusion through bovine dentine, 622
bovine teeth, Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267

calcium chelation, Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
calcium hydroxide, Chemical and antimicrobial properties of calcium hydroxide mixed with irrigating solutions, 100
Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267
Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
Mineral trioxide aggregate (MTA) and calcium hydroxide as pulp-capping agents in human teeth: a preliminary report, 225
Remaining dentine thickness and human pulp responses, 33
The relationship of intracanal medicaments to post-operative pain in endodontics, 868
Campylobacter gracilis, *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections, 174
Campylobacter rectus, *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections, 174
canal cleanliness, Light and scanning electron microscopic evaluation of Glyde™ File Prep in smear layer removal, 336
canal curvature, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
canal filling, Comparison of MicroSeal and System B/Obtura II obturation techniques, 876
canal geometry, ProTaper rotary root canal preparation: effects of canal anatomy on final shape analysed by micro CT, 86
canal irrigants, Comparison of antibacterial and toxic effects of various root canal irrigants, 423
canal preparation, A combined radiographic and computerized scanning method to evaluate remaining dentine thickness in mandibular incisors after various intracanal procedures, 682
canal shape, Root-canal preparation with FlexMaster: canal shapes analysed by micro-computed tomography, 740
canal shaping, Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
canal straightening, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
canal transportation, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
canal wall thickness, A comparison of the minimum canal wall thickness remaining following preparation using two nickel–titanium rotary systems, 636
Candida albicans, Examination on *Candida* spp. in refractory periapical granulomas, 643
caries-affected dentine, Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
caries, Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64
Carisolv™, Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
case classification, An evaluation of the usefulness of two endodontic case assessment forms by general dentists, 545
case report, Orthograde retreatment and apexification after unsuccessful endodontic treatment, retreatment and apicectomy, 780
catalase, Catalase activity in human healthy and inflamed dental pulps, 599
cavity preparation, Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106

© 2003 Blackwell Publishing Ltd

CDJ. The location of the cementodentinal junction in single-rooted mandibular first premolars from Egyptian and Saudi patients: a histological study, 541
cell culture. Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404
cementoma. Focal cemento-osseous dysplasia involving a mandibular lateral incisor, 907
cetrimide. Comparison of antibacterial and toxic effects of various root canal irrigants, 423
cGMP. Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
CGRP. Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
chelators. Chelating agents in root canal treatment: mode of action and indications for their use, 810
chlorhexidine gluconate. Comparison of antibacterial and toxic effects of various root canal irrigants, 423
chlorhexidine. Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267
Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
Scanning electron microscopic study of the cleaning ability of chlorhexidine as a root-canal irrigant, 391
The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine *in vitro*, 848
chromosomal aberrations. Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
cleaning procedures. The effect of cleaning procedures on fracture properties and corrosion of NiTi files, 724
cleaning. Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
clinical follow up. Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787
combinations. Chemical and antimicrobial properties of calcium hydroxide mixed with irrigating solutions, 100
compomer. Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281
composite. Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281
composition. Chemical and X-ray analyses of five brands of dental gutta-percha cone, 302
computerized photography. A combined radiographic and computerized scanning method to evaluate remaining dentine thickness in mandibular incisors after various intracanal procedures, 682
coronal leakage. Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787
corrosion. The effect of cleaning procedures on fracture properties and corrosion of NiTi files, 724
corticosteroids. The relationship of intracanal medicaments to postoperative pain in endodontics, 868
current-clamp. Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
curved root canals. Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
Preparation of curved root canals with rotary FlexMaster instruments compared to Lightspeed instruments and NiTi hand files, 483
cyclic fatigue. ProTaper rotary root canal preparation: assessment of torque and force in relation to canal anatomy, 93
Root canal preparation with FlexMaster: asessment of torque and force in relation to canal anatomy, 883

debris. Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
decision making. Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
degradation. Solubility of root-canal sealers in water and artificial saliva, 660
demineralization. A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
dental cone. Chemical and X-ray analyses of five brands of dental gutta-percha cone, 302
dental education. Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
dental materials. Rheological properties and biocompatibility of endodontic sealers, 527
dental practice. Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
dental pulp chamber. Coronal pulp size in molars: a study of bitewing radiographs, 757

© 2003 Blackwell Publishing Ltd

dental pulp. Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
Catalase activity in human healthy and inflamed dental pulps, 599
Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404
Lymphangiogenesis in human dental pulp, 853
dental referral. Factors influencing referral for specialist endodontic treatment amongst a group of Dutch general practitioners, 129
dentine bonding. *In vitro* fracture resistance of root-filled teeth using new-generation dentine bonding adhesives, 770
dentine hardness. Chelating agents in root canal treatment: mode of action and indications for their use, 810
dentine permeability. Chelating agents in root canal treatment: mode of action and indications for their use, 810
difficulty. An evaluation of the usefulness of two endodontic case assessment forms by general dentists, 545
dimethylformamide. Removal of gutta-percha from root canals using an Nd:YAG laser, 670
disintegration. Solubility of root-canal sealers in water and artificial saliva, 660
double teeth. Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372
dye penetration. Evaluation of root-end cavity preparation using ultrasonic retrotips, 586

EBA. Investigation of the marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491
echography. Ultrasound real-time imaging in the differential diagnosis of periapical lesions, 556
EDTA. Chelating agents in root canal treatment: mode of action and indications for their use, 810
Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
education. Teaching an engine-driven preparation technique to undergraduates: initial observations, 476
endodontic cements. A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
Endodontic Cube. A comparison of the minimum canal wall thickness remaining following preparation using two nickel–titanium rotary systems, 636
endodontic diagnosis. Ultrasound real-time imaging in the differential diagnosis of periapical lesions, 556
endodontic failure. Microorganisms from canals of root-filled teeth with periapical lesions, 1
endodontic infection. Oral treponemes in primary root canal infections as detected by nested PCR, 20
endodontic infections. *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections, 174
endodontic materials. Short-term periradicular tissue response to mineral trioxide aggregate (MTA) as root-end filling material, 44
endodontic radiography. Effect of X-ray beam angulation and intraradicular contrast medium on radiographic interpretation of lower first molar root canal anatomy, 12
endodontic sealer. Comparative *in vivo* analysis of the sealing ability of three endodontic sealers in post-prepared root canals, 857
endodontic sealers. Solubility of root-canal sealers in water and artificial saliva, 660
endodontic surgery. A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
endodontic therapy. The relationship of intracanal medicaments to postoperative pain in endodontics, 868
endodontic treatment. An evaluation of the usefulness of two endodontic case assessment forms by general dentists, 545
Diagnosis and root canal treatment of a mandibular second premolar with three root canals, 912
Endodontic and surgical treatment of a geminated maxillary incisor, 380
Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372
Microbial causes of endodontic flare-ups, 453
endodontics. *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical gutta-percha condensation techniques, 367
A preliminary *in vitro* study of the incidence and position of the root canal isthmus in maxillary and mandibular first molars, 276
Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787
Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 1. Intracanal drugs and substances, 75
Biocompatibility of dental materials used in contemporary endodontic therapy: a review. Part 2. Root-canal-filling materials, 147
Coronal pulp size in molars: a study of bitewing radiographs, 757
Effect of Er:YAG laser and EDTAC on the adhesiveness to dentine of different sealers containing calcium hydroxide, 472
Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations, 181

© 2003 Blackwell Publishing Ltd

Rheological properties and biocompatibility of endodontic sealers, 527
endoscopy, Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
endothelium-dependent vasodilatation, Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
endothelium, Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
endotoxin, Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
epidemiology, Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations, 181
Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
Er:YAG laser, Effect of Er:YAG laser and EDTAC on the adhesiveness to dentine of different sealers containing calcium hydroxide, 472
eucalyptol, Removal of gutta-percha from root canals using an Nd:YAG laser, 670

facial radicular groove, Bilateral buccal radicular groove in maxillary incisors: case report, 898
failure, Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
fibre optic orascope, Orascopic visualization technique for conventional and surgical endodontics, 441
fibro-osseous lesions, Focal cemento-osseous dysplasia involving a mandibular lateral incisor, 907
filling, Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591
flap design, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
flare-up, Microbial causes of endodontic flare-ups, 453
FlexMaster, A comparative study of root canal preparation using FlexMaster and HERO 642 rotary Ni–Ti instruments, 358
Root canal preparation with FlexMaster: asessment of torque and force in relation to canal anatomy, 883
Root-canal preparation with FlexMaster: canal shapes analysed by micro-computed tomography, 740
flow, Rheological properties and biocompatibility of endodontic sealers, 527
force, ProTaper rotary root canal preparation: assessment of torque and force in relation to canal anatomy, 93
Root canal preparation with FlexMaster: asessment of torque and force in relation to canal anatomy, 883
fracture properties, The effect of cleaning procedures on fracture properties and corrosion of NiTi files, 724
fracture resistance, *In vitro* fracture resistance of root-filled teeth using new-generation dentine bonding adhesives, 770
fracture, Factors influencing the fracture of nickel-titanium rotary instruments, 262
furcal accessory canal incidence, Incidence of patent furcal accessory canals in permanent molars of a Turkish population, 515
fused tooth, Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
fusion, Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372

gemination, Endodontic and surgical treatment of a geminated maxillary incisor, 380
Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372
gingival fibroblasts, Effect of apical root-end filling materials on gingival fibroblasts, 610
glass-ionomer cement, Acidity of conventional luting cements and their diffusion through bovine dentine, 622
Biocompatibility of resin-based materials used as pulp-capping agents, 831
gutta-percha, A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
Chemical and X-ray analyses of five brands of dental gutta-percha cone, 302
Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591
The antimicrobial efficacy of 'MGP' gutta-percha *in vitro*, 616

Helix neurones, Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
hemisection, Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
Endodontic and surgical treatment of a geminated maxillary incisor, 380
HERO 642, A comparative study of root canal preparation using FlexMaster and HERO 642 rotary Ni–Ti instruments, 358
Herofill™ Soft-core obturation, *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical gutta-percha condensation techniques, 367
histologic evaluation, Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64
human lymphocytes, Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330

© 2003 Blackwell Publishing Ltd

hydrogen peroxide, Catalase activity in human healthy and inflamed dental pulps, 599
Review of the current status of tooth whitening with the walking bleach technique, 313
hyperplastic pulpitis, Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64

IL-6, Induction of interleukin-6 gene expression by pro-inflammatory cytokines and black-pigmented *Bacteroides* in human pulp cell cultures, 352
IL-8, Induction of interleukin-8 gene expression by black-pigmented *Bacteroides* in human pulp fibroblasts and osteoblasts, 774
immunohistochemical methods, Lymphangiogenesis in human dental pulp, 853
immunohistochemistry, Inflammatory infiltrate of chronic periradicular lesions: an immunohistochemical study, 464
in vitro, The antimicrobial efficacy of 'MGP' gutta-percha *in vitro*, 616
incidence, A preliminary *in vitro* study of the incidence and position of the root canal isthmus in maxillary and mandibular first molars, 276
incisor, Focal cemento-osseous dysplasia involving a mandibular lateral incisor, 907
infiltrate, Inflammatory infiltrate of chronic periradicular lesions: an immunohistochemical study, 464
inflammation, Catalase activity in human healthy and inflamed dental pulps, 599
Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106
Remaining dentine thickness and human pulp responses, 33
injury, Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106
inner layer of dentine, The capability of two hand instrumentation techniques to remove the inner layer of dentine in oval canals, 218
instrumentation, The capability of two hand instrumentation techniques to remove the inner layer of dentine in oval canals, 218
intact dentine, Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
intracanal medicament, Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
intracanal medicaments, Effectiveness of 2% chlorhexidine gel and calcium hydroxide against *Enterococcus faecalis* in bovine root dentine *in vitro*, 267
The relationship of intracanal medicaments to post-operative pain in endodontics, 868
iodoform, The antimicrobial efficacy of 'MGP' gutta-percha *in vitro*, 616
iontophoresis, Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674

IRM, A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine, 604
irrigant, Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
irrigants, Scanning electron microscopic study of the cleaning ability of chlorhexidine as a root-canal irrigant, 391
The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine *in vitro*, 848
irrigation, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
isthmus, A preliminary *in vitro* study of the incidence and position of the root canal isthmus in maxillary and mandibular first molars, 276

K3, An *in vitro* study of the torsional properties of new and used K3 instruments, 764

L-NAME, Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
leakage, Comparative *in vivo* analysis of the sealing ability of three endodontic sealers in post-prepared root canals, 857
Fluid movement along the coronal two-thirds of root fillings placed by three different gutta-percha techniques, 533
Ledermix, The relationship of intracanal medicaments to postoperative pain in endodontics, 868
Lightspeed, Comparative study of root-canal preparation using Lightspeed and Quantec SC rotary NiTi instruments, 748
Comparison of procedural errors resulting during root canal preparations completed by junior dental students in patients using an '8-step method' versus 'serial step-back technique', 49
long-term results, Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
longevity, Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
lymphangiogenesis, Lymphangiogenesis in human dental pulp, 853

mandibular first premolar, The location of the cemento-dentinal junction in single-rooted mandibular first premolars from Egyptian and Saudi patients: a histological study, 541
mandibular incisors, A combined radiographic and computerized scanning method to evaluate remaining

© 2003 Blackwell Publishing Ltd

dentine thickness in mandibular incisors after various intracanal procedures, 682
mandibular second premolar, Diagnosis and root canal treatment of a mandibular second premolar with three root canals, 912
marginal adaptation, Investigation of the marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491
maxillary lateral incisor, Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
maxillary premolars, Root canal treatment of three-rooted maxillary first and second premolars – a case report, 705
metronidazole, Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
MG-63 osteosarcoma cells, Osteoblasts and MG-63 osteosarcoma cells behave differently when in contact with ProRoot™ MTA and White MTA, 564
'MGP' gutta-percha, The antimicrobial efficacy of 'MGP' gutta-percha *in vitro*, 616
micro-computed tomography (μCT), Root-canal preparation with FlexMaster: canal shapes analysed by micro-computed tomography, 740
microbiology, Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
Microorganisms from canals of root-filled teeth with periapical lesions, 1
microcrack, Evaluation of root-end cavity preparation using ultrasonic retrotips, 586
micronucleus, Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
mineral trioxide aggregate (MTA), Mineral trioxide aggregate (MTA) and calcium hydroxide as pulp-capping agents in human teeth: a preliminary report, 225
mineral trioxide aggregate, Osteoblasts and MG-63 osteosarcoma cells behave differently when in contact with ProRoot™ MTA and White MTA, 564
Short-term periradicular tissue response to mineral trioxide aggregate (MTA) as root-end filling material, 44
molar morphology, Incidence of patent furcal accessory canals in permanent molars of a Turkish population, 515
MTA, A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
Effect of apical root-end filling materials on gingival fibroblasts, 610
Investigation of the marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491

NaOCl, Comparison of antibacterial and toxic effects of various root canal irrigants, 423
naproxen sodium, Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
Nd:YAG laser, Removal of gutta-percha from root canals using an Nd:YAG laser, 670
nested PCR, *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections, 174
neurotoxicity, Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
Ni–Ti instruments, A comparative study of root canal preparation using FlexMaster and HERO 642 rotary Ni–Ti instruments, 358
nickel–titanium, A comparison of the minimum canal wall thickness remaining following preparation using two nickel–titanium rotary systems, 636
Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
Incidence of rotary ProFile instrument fracture and the potential for bypassing *in vivo*, 864
Preparation of curved root canals with rotary FlexMaster instruments compared to Lightspeed instruments and NiTi hand files, 483
Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
nickel-titanium, Factors influencing the fracture of nickel-titanium rotary instruments, 262
Influence of operator experience on canal preparation time when using the rotary Ni-Ti ProFile system in simulated curved canals, 161
Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
NIT, Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591
NiTi files, Comparison of procedural errors resulting during root canal preparations completed by junior dental students in patients using an '8-step method' versus 'serial step-back technique', 49
The effect of cleaning procedures on fracture properties and corrosion of NiTi files, 724
NiTi instruments, Comparative study of root-canal preparation using Lightspeed and Quantec SC rotary NiTi instruments, 748
nitric oxide, Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840

obturation, A systematic review of *in vivo* retrograde obturation materials, 577
Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
occlusal load, Measurement of periapical pressure created by occlusal loading, 700
odontoblast, Remaining dentine thickness and human pulp responses, 33

© 2003 Blackwell Publishing Ltd

oral spirochaetes, Oral treponemes in primary root canal infections as detected by nested PCR, 20
orascopic endodontics, Orascopic visualization technique for conventional and surgical endodontics, 441
orthograde, Orthograde retreatment and apexification after unsuccessful endodontic treatment, retreatment and apicectomy, 780
osteoblasts, Induction of interleukin-8 gene expression by black-pigmented *Bacteroides* in human pulp fibroblasts and osteoblasts, 774
Osteoblasts and MG-63 osteosarcoma cells behave differently when in contact with ProRoot™ MTA and White MTA, 564
osteonectin, Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404

pain, The relationship of intracanal medicaments to postoperative pain in endodontics, 868
papilla base incision, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
papilla healing, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
patent furcal accessory canals, Incidence of patent furcal accessory canals in permanent molars of a Turkish population, 515
periapical disease, Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787
periapical diseases, Focal cemento-osseous dysplasia involving a mandibular lateral incisor, 907
periapical granuloma, Examination on *Candida* spp. in refractory periapical granulomas, 643
periapical lesions, Ultrasound real-time imaging in the differential diagnosis of periapical lesions, 556
periodontal regeneration, Bilateral buccal radicular groove in maxillary incisors: case report, 898
periodontal–endodontic lesions, Bilateral buccal radicular groove in maxillary incisors: case report, 898
periradicular lesions, Inflammatory infiltrate of chronic periradicular lesions: an immunohistochemical study, 464
periradicular pathosis, Short-term periradicular tissue response to mineral trioxide aggregate (MTA) as root-end filling material, 44
periradicular surgery, Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
pH, Acidity of conventional luting cements and their diffusion through bovine dentine, 622
Chemical and antimicrobial properties of calcium hydroxide mixed with irrigating solutions, 100
physical properties, Solubility of root-canal sealers in water and artificial saliva, 660
polymerase chain reaction, Oral treponemes in primary root canal infections as detected by nested PCR, 20
preparation time, Evaluation of root-end cavity preparation using ultrasonic retrotips, 586
preparation, Teaching an engine-driven preparation technique to undergraduates: initial observations, 476
pro-inflammatory cytokines, Induction of interleukin-6 gene expression by pro-inflammatory cytokines and black-pigmented *Bacteroides* in human pulp cell cultures, 352
ProFile, Incidence of rotary ProFile instrument fracture and the potential for bypassing *in vivo*, 864
profile, Comparison of procedural errors resulting during root canal preparations completed by junior dental students in patients using an '8-step method' versus 'serial step-back technique', 49
properties, A comparative study of selected physical properties of five root-canal sealers, 629
prospective study, Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
prostaglandin release, Effect of apical root-end filling materials on gingival fibroblasts, 610
ProTaper, ProTaper rotary root canal preparation: assessment of torque and force in relation to canal anatomy, 93
ProTaper rotary root canal preparation: effects of canal anatomy on final shape analysed by micro CT, 86
protein synthesis, Effect of apical root-end filling materials on gingival fibroblasts, 610
pulp cap, Mineral trioxide aggregate (MTA) and calcium hydroxide as pulp-capping agents in human teeth: a preliminary report, 225
pulp fibroblasts, Induction of interleukin-6 gene expression by pro-inflammatory cytokines and black-pigmented *Bacteroides* in human pulp cell cultures, 352
Induction of interleukin-8 gene expression by black-pigmented *Bacteroides* in human pulp fibroblasts and osteoblasts, 774
pulp, Biocompatibility of resin-based materials used as pulp-capping agents, 831
pulpal infection, Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500

quality of filling, Comparison of MicroSeal and System B/Obtura II obturation techniques, 876
Quantec SC, Comparative study of root-canal preparation using Lightspeed and Quantec SC rotary NiTi instruments, 748

radiographic apex, The location of the cementodentinal junction in single-rooted mandibular first premolars from Egyptian and Saudi patients: a histological study, 541

© 2003 Blackwell Publishing Ltd

radiography, Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
radiology, Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations, 181
radiopaque contrast medium, Effect of X-ray beam angulation and intraradicular contrast medium on radiographic interpretation of lower first molar root canal anatomy, 12
recession, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
referral, Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
regeneration, Histomorphometric analysis of odontoblast-like cell numbers and dentine bridge secretory activity following pulp exposure, 106
remaining dentine thickness., A combined radiographic and computerized scanning method to evaluate remaining dentine thickness in mandibular incisors after various intracanal procedures, 682
remaining dentine thickness, Acidity of conventional luting cements and their diffusion through bovine dentine, 622
reparative dentine, Remaining dentine thickness and human pulp responses, 33
resin blocks, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
retreatment, Orthograde retreatment and apexification after unsuccessful endodontic treatment. retreatment and apicectomy, 780
Removal of gutta-percha from root canals using an Nd:YAG laser, 670
retrograde, A systematic review of *in vivo* retrograde obturation materials, 577
rod-lens endoscope, Orascopic visualization technique for conventional and surgical endodontics, 441
root anatomy, Root canal treatment of three-rooted maxillary first and second premolars – a case report, 705
root canal anatomy, Effect of X-ray beam angulation and intraradicular contrast medium on radiographic interpretation of lower first molar root canal anatomy, 12
root canal conditioner, Light and scanning electron microscopic evaluation of Glyde™ File Prep in smear layer removal, 336
root canal filling, Comparative *in vivo* analysis of the sealing ability of three endodontic sealers in post-prepared root canals, 857
Tissue reaction initiated by different sealers, 256
root canal infection, Microbial causes of endodontic flare-ups, 453
root canal preparation, Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
Influence of operator experience on canal preparation time when using the rotary Ni-Ti ProFile system in simulated curved canals, 161
Preparation of curved root canals with rotary FlexMaster instruments compared to Lightspeed instruments and NiTi hand files, 483
Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
root canal sealers, Effect of Er:YAG laser and EDTAC on the adhesiveness to dentine of different sealers containing calcium hydroxide, 472
root canal treatment, Chelating agents in root canal treatment: mode of action and indications for their use, 810
Comparison of procedural errors resulting during root canal preparations completed by junior dental students in patients using an '8-step method' versus 'serial step-back technique', 49
Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
Radiographic features and outcome of root canal treatment carried out in the Łódź region of Poland, 27
Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
Technical quality of root canal treatment in Taiwan, 416
root canal, A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
root resorption, Endodontic and orthodontic treatment of a cross-bite fused maxillary lateral incisor, 135
root-canal filling, *In vitro* evaluation of apical microleakage following canal filling with a coated carrier system compared with lateral and thermomechanical gutta-percha condensation techniques, 367
root-canal infection, Examination on *Candida* spp. in refractory periapical granulomas, 643
root-canal posts, Factors influencing the removal of posts, 687
root-canal preparation, Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
root-canal sealer, Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide–eugenol-based sealer, 296
root-canal therapy, Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment, 500
Microorganisms from canals of root-filled teeth with periapical lesions, 1
root-canal treatment, Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases, 787

© 2003 Blackwell Publishing Ltd

Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
root-end cavity preparation. Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
root-end cavity. Evaluation of root-end cavity preparation using ultrasonic retrotips, 586
root-end filling. A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
Investigation of the marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491
Short-term periradicular tissue response to mineral trioxide aggregate (MTA) as root-end filling material, 44
root-end resection. Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
root-filled teeth. *In vitro* fracture resistance of root-filled teeth using new-generation dentine bonding adhesives, 770
rotary files. Influence of operator experience on canal preparation time when using the rotary Ni-Ti ProFile system in simulated curved canals, 161
Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
rotary instrument. Preparation of curved root canals with rotary FlexMaster instruments compared to Lightspeed instruments and NiTi hand files, 483
rotary instruments. Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 1. Shaping ability in simulated curved canals, 199
Factors influencing the fracture of nickel-titanium rotary instruments, 262
rotary. Teaching an engine-driven preparation technique to undergraduates: initial observations, 476
rotational speed. Factors influencing the fracture of nickel-titanium rotary instruments, 262

sealer. Fluid movement along the coronal two-thirds of root fillings placed by three different gutta-percha techniques, 533
sealers. A comparative study of selected physical properties of five root-canal sealers, 629
Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
Tissue reaction initiated by different sealers, 256
SEM. Diagnostic accuracy of endoscopy in periradicular surgery – a comparison with scanning electron microscopy, 691
Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
separation. Incidence of rotary ProFile instrument fracture and the potential for bypassing *in vivo*, 864
setting time. Fluid movement along the coronal two-thirds of root fillings placed by three different gutta-percha techniques, 533
shaft design. Progressive versus constant tapered shaft design using NiTi rotary instruments, 288
shape. ProTaper rotary root canal preparation: effects of canal anatomy on final shape analysed by micro CT, 86
shaping. Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
silicone. Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide–eugenol-based sealer, 296
simulated canals. Influence of operator experience on canal preparation time when using the rotary Ni-Ti ProFile system in simulated curved canals, 161
simulated root canals. Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
smear layer. Chelating agents in root canal treatment: mode of action and indications for their use, 810
Efficiency of rotary nickel–titanium K3 instruments compared with stainless steel hand K-Flexofile. Part 2. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth, 208
Light and scanning electron microscopic evaluation of Glyde™ File Prep in smear layer removal, 336
Scanning electron microscopic study of the cleaning ability of chlorhexidine as a root-canal irrigant, 391
sodium hypochlorite. Effect of different irrigation solutions and calcium hydroxide on bacterial LPS, 733
Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine *in vitro*, 848
sodium perborate. Review of the current status of tooth whitening with the walking bleach technique, 313
sodium salicylate. Facilitation of iontophoretic drug delivery through intact and caries-affected dentine, 674
specialist endodontic treatment. Factors influencing referral for specialist endodontic treatment amongst a group of Dutch general practitioners, 129
spontaneous activity. Comparison of neurotoxicity of root canal sealers on spontaneous bioelectrical activity in identified *Helix* neurones using an intracellular recording technique, 891
subcutaneous connective tissue. Reactions of connective tissue to compomers, composite and amalgam root-end filling materials, 281

© 2003 Blackwell Publishing Ltd

Substance P, Effect of nitric oxide synthase inhibitor (L-NAME) on substance P-induced vasodilatation in the dental pulp, 840
success rate, A prospective clinical study of Mineral Trioxide Aggregate and IRM when used as root-end filling materials in endodontic surgery, 520
success, Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
sulcular incision, Comparison of papilla healing following sulcular full-thickness flap and papilla base flap in endodontic surgery, 653
supernumerary teeth, Endodontic treatment of developmental anomalies in posterior teeth: treatment of geminated/fused teeth – report of two cases, 372
surgical endodontics, Periapical endodontic surgery: a 3-year follow-up study, 193
survey, Endodontic treatment performed by Flemish dentists. Part 2. Canal filling and decision making for referrals and treatment of apical periodontitis, 344
Root canal treatment performed by Flemish dentists. Part 1. Cleaning and shaping, 166
survival analysis, Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
systematic review, A systematic review of *in vivo* retrograde obturation materials, 577

Taiwan, Technical quality of root canal treatment in Taiwan, 416
teaching, Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
technical quality, Technical quality of root canal treatment in Taiwan, 416
thermomechanical, Comparison of MicroSeal and System B/Obtura II obturation techniques, 876
tissue dissolution, Interactions of ethylenediamine tetraacetic acid with sodium hypochlorite in aqueous solutions, 411
tissue reaction, Tissue reaction initiated by different sealers, 256
titanium, Effect of apical root-end filling materials on gingival fibroblasts, 610
tooth clearing, A preliminary analysis of the morphology of lateral canals after root canal filling using a tooth-clearing technique, 54
tooth resorption, Review of the current status of tooth whitening with the walking bleach technique, 313
torque at fracture, An *in vitro* study of the torsional properties of new and used K3 instruments, 764
torque, ProTaper rotary root canal preparation: assessment of torque and force in relation to canal anatomy, 93
Root canal preparation with FlexMaster: asessment of torque and force in relation to canal anatomy, 883
toxic effects, Comparison of antibacterial and toxic effects of various root canal irrigants, 423
transportation, ProTaper rotary root canal preparation: effects of canal anatomy on final shape analysed by micro CT, 86
Root-canal preparation with FlexMaster: canal shapes analysed by micro-computed tomography, 740
trauma, Histological evaluation of teeth with hyperplastic pulpitis caused by trauma or caries: case reports, 64
treatment outcome, Long-term survival of primary root canal treatment carried out in a dental teaching hospital, 117
Turkish subpopulation, Incidence of patent furcal accessory canals in permanent molars of a Turkish population, 515
type I collagen, Immunolocalization of bone extracellular matrix proteins (type I collagen, osteonectin and bone sialoprotein) in human dental pulp and cultured pulp cells, 404

ultrasonic preparation, Investigation of the marginal adaptation of root-end filling materials in root-end cavities prepared with ultrasonic tips, 491
ultrasonic retrotip, Evaluation of root-end cavity preparation using ultrasonic retrotips, 586
ultrasound, Factors influencing the removal of posts, 687
Periapical endodontic surgery: a 3-year follow-up study, 193
Preliminary *in vitro* evaluation of Carisolv™ as a root canal irrigant, 433
ultrastructure, Age-related changes in blood capillary endothelium of human dental pulp: an ultrastructural study, 395
undergraduate, Root canal shaping with manual stainless steel files and rotary Ni–Ti files performed by students, 246
Root-canal shaping with manual and rotary Ni-Ti files performed by students, 715
Teaching an engine-driven preparation technique to undergraduates: initial observations, 476

V79, Examination of cytotoxicity and mutagenicity of AH26 and AH Plus sealers, 330
VEGFR-3, Lymphangiogenesis in human dental pulp, 853
vertical compaction, Percentage of filled canal area in mandibular molars after conventional root-canal instrumentation and after a noninstrumentation technique (NIT), 591

weight loss, Solubility of root-canal sealers in water and artificial saliva, 660

XMCT, Progressive versus constant tapered shaft design using NiTi rotary instruments, 288

© 2003 Blackwell Publishing Ltd

yeasts. Examination on *Candida* spp. in refractory periapical granulomas, 643

zinc oxide eugenol. Remaining dentine thickness and human pulp responses, 33
zinc phosphate cement. Acidity of conventional luting cements and their diffusion through bovine dentine, 622
zinc polycarboxylate cement. Acidity of conventional luting cements and their diffusion through bovine dentine, 622
ZOE. Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide–eugenol-based sealer, 296

© 2003 Blackwell Publishing Ltd